

Non-equilibrium relaxation of an elastic string in random media

Alejandro B. Kolton¹, Alberto Rosso² and Thierry Giamarchi¹

¹ *Université de Genève, DPMC, 24 Quai Ernest Ansermet, CH-1211 Genève 4, Switzerland.*

² *Laboratoire de physique théorique et modèles statistiques CNRS UMR8626, Bât. 100 Université Paris-Sud; 91405 Orsay Cedex, France.*

Abstract

We study the relaxation of an elastic string in a two dimensional pinning landscape using Langevin dynamics simulations. The relaxation of a line, initially flat, is characterized by a growing length, $L(t)$, separating the equilibrated short length scales from the flat long distance geometry that keep memory of the initial condition. We find that, in the long time limit, $L(t)$ has a non-algebraic growth, consistent with thermally activated jumps over barriers with power law scaling, $U(L) \sim L^\theta$.

Understanding the dynamics of elastic manifolds in random media has been the focus of intense activities both on the theoretical and experimental side [1]. Because of the competition between disorder and elasticity, glassy properties arise leading to divergent barriers separating metastable states. The creep, *i.e.* the steady-state response to a small external force, has been studied theoretically [2,3] and observed experimentally [4]. Much less is known about the non-stationary relaxation to equilibrium. Theoretical attempts to tackle this problem have been made using mean field and renormalization group approaches [5,6], while numerical simulations have been mostly restricted to 2-dimensional systems [7]. Direct applications of these results to one dimensional domain walls are however difficult.

In this paper we study, numerically, the slow non-stationary motion of an elastic string in a two dimensional random medium, relaxing from a flat initial configuration. The string obeys the equation of motion $\gamma \partial_t u(z, t) = c \partial_z^2 u(z, t) + F_p(u, z) + \eta(z, t)$, where γ is the friction coefficient and c the elastic constant, $F_p(u, z)$ the pinning force for a random bond disorder and $\eta(z, t)$ is the thermal noise. The details of the numerical method are given in [8].

Fig. 1(a) shows the typical relaxation of a string and Fig. 1(b) the evolution of the structure factor $S(q, t)$ which characterize the geometry of the line [3,8]. At short length scales the line has reached equilibrium and it is characterized by the well known roughness exponent $\zeta = 2/3$ [9]. At large length scales a plateau still keeps memory of the initial flat condition. A unique crossover length $L(t)$ can be defined. Its evolution is shown in Fig. 1(c). Two main scenarios have been proposed to describe the long time growth of $L(t)$. The first one relies on phenomenological scaling arguments, based on creep. At low temperatures the relaxation is dominated by the energy barriers $U(L)$ that must be overcome in order to equilibrate the system up to a length scale L . Using the Arrhenius thermal activation law we can thus express the relaxation time $t(L) \sim \exp[\beta U(L)]$. Even if the exact numerical determination of $U(L)$ is an NP-complete problem it is usually conjectured that the typical barriers of the energy landscape scale, asymptotically with L , the same way as the free energy fluctuations:




Fig. 1. (a) Typical configurations of the relaxing string (b) $S(q, t)$ at $T = 0.5$. The dashed line corresponds thermal equilibrium solution. (c) Characteristic length scale $L(t)$ of a string of size $L = 256$. ($\diamond$) symbols and the dotted line correspond to the numerical and analytical $L(t)$ for the clean system, and ($\bullet$) symbols to the disordered case. The continuum line fits to the logarithmic growth, and the dashed line is a fit to a power law growth at intermediate scales. Inset: exponent θ of the logarithmic law extracted from the fit in the time interval $t_i < t < 10^6$, for $L = 512$ ($\triangle$) and $L = 256$ ($\circ$).

$U(L) \sim L^\theta$, with $\theta = 1/3$ for a line. Numerical calculations [11] and FRG calculations [2] seem to confirm this conjecture. Following these arguments we infer that $L(t) \sim [\log(t)]^{1/\theta}$ [8, 10]. The second scenario predicts $L(t) \sim t^{1/z}$, like for the clean system but with a new temperature dependent exponent $z > 2$ [6, 7]. This power law scaling can be ruled out from our data, due to the observed bending in the log-log scale in Fig. 1(c). On the other hand a two-parameters fit to the logarithmic growth law gives an exponent θ which, at long times, becomes size and time independent, as shown in the inset of Fig. 1. However we found $\theta \approx 0.49$. This value, bigger than the expected $\theta = 1/3$, indicates either a violation of the expected scaling of barriers or the presence of logarithmic corrections in the leading term: $U(L) \sim L^{1/3} \log^\mu(L)$ [10, 11].

References

- [1] T. Giamarchi, A. B. Kolton, and A. Rosso, Proceedings of the XIX Sitges conference on “Jamming, Yielding, and Irreversible Deformation in Condensed Matter”, Sitges, Spain (2004), [cont-mat/0503437](#).
- [2] L. B. Ioffe and V. M. Vinokur, J. Phys. C **20**, 6149 (1987); T. Nattermann, Europhys. Lett. **4**, 1241 (1987); P. Chauve, T. Giamarchi, and P. Le Doussal, Phys. Rev. B **62**, 6241 (2000).
- [3] A. B. Kolton, A. Rosso, and T. Giamarchi, Phys. Rev. Lett. **94** 047002 (2005).
- [4] S. Lemerle *et al.*, Phys. Rev. Lett. **80**, 849 (1998); T. Tybell *et al.*, Phys. Rev. Lett. **89**, 097601 (2002).
- [5] L. Cugliandolo, J. Kurchan, and P. Le Doussal, Phys. Rev. Lett. **76**, 2390 (1996). L. Balents and P. Le Doussal, Phys. Rev. E **69**, 061107 (2004);
- [6] G. Schehr and P. Le Doussal, Phys. Rev. Lett. **93**, 217201 (2004); *ibid* [cond-mat/0501199](#), 2005.
- [7] G. Schehr and H. Rieger, Phys. Rev. B **71**, 184202 (2005); H. Rieger, G. Schehr, and R. Paul, [cond-mat/0411234](#), 2004.
- [8] A. B. Kolton, A. Rosso, and T. Giamarchi, Phys. Rev. Lett. **95**, 180604 (2005)
- [9] M. Kardar, Phys. Rev. Lett. **55**, 2923 (1985).
- [10] H. Yoshino, private communication, unpublished.
- [11] B. Drossel and M. Kardar, Phys. Rev. E **52**, 4841 (1995).